

# Measure Theory with Ergodic Horizons

## Lecture 2

### Generation of algebras and $\sigma$ -algebras.

Let  $X$  be a set and  $\mathcal{C} \subseteq \mathcal{P}(X)$  be a collection. Is there a smallest  $(\sigma)$ -algebra containing  $\mathcal{C}$ ? There is at least one  $(\sigma)$ -algebra containing  $\mathcal{C}$ , namely,  $\mathcal{P}(X)$ .

Observation. Arbitrary intersections of  $(\sigma)$ -algebras is still a  $(\sigma)$ -algebra, i.e. if  $\{\mathcal{A}_i : i \in I\}$  is a collection of  $(\sigma)$ -algebras on  $X$ , then  $\bigcap_{i \in I} \mathcal{A}_i$  is a  $(\sigma)$ -algebra.

Thus, given  $\mathcal{C} \subseteq \mathcal{P}(X)$ ,  $\langle \mathcal{C} \rangle :=$  intersection of all algebras containing  $\mathcal{C}$   
 $\langle \mathcal{C} \rangle_{\sigma} :=$  intersection of all  $\sigma$ -algebras containing  $\mathcal{C}$   
is the smallest  $(\sigma)$ -algebra containing  $\mathcal{C}$ .

These are top-down definitions, which are hard to work with, so we seek bottom-up definitions.

Prop. Let  $\mathcal{C} \subseteq \mathcal{P}(X)$ . Then

(a)  $\langle \mathcal{C} \rangle = \bigcup_{n \in \mathbb{N}} \mathcal{C}_n$ , where  $\mathcal{C}_0 := \mathcal{C} \cup \{\emptyset\}$ ,  $\mathcal{C}_{n+1} := \{C^c : C \in \mathcal{C}_n\} \cup \{\text{finite unions of sets in } \mathcal{C}_n\}$ .

(b)  $\langle \mathcal{C} \rangle_{\sigma} = \bigcup_{\alpha < \omega_1} \mathcal{C}_{\alpha}$ , where  $\mathcal{C}_0 := \mathcal{C} \cup \{\emptyset\}$ ,  $\mathcal{C}_{\alpha} := \bigcup_{\beta < \alpha} \{C^c : C \in \mathcal{C}_{\beta}\} \cup \{\text{ctbl unions of sets in } \bigcup_{\beta < \alpha} \mathcal{C}_{\beta}\}$ .  
 $\alpha < \omega_1$  ← first unctbl ordinal

Proof. (a) is left as HW and (b) is an optional exercise.  $\square$

Def. For a metric (more generally, topological) space  $X$ , the  $\sigma$ -algebra generated by all open sets is called the **Borel  $\sigma$ -algebra**, and the sets in it are called **Borel sets**. We denote the Borel  $\sigma$ -algebra of  $X$  by  $\mathcal{B}(X)$ .

Observe that the Borel  $\sigma$ -algebra is also generated by closed sets.

Def. A **basis** for a metric (topological space)  $X$  is a collection  $\mathcal{C}$  of open sets such that each open set in  $X$  is a union of some sets from  $\mathcal{C}$ .  
 $X$  is called **second ctbl** if it admits a ctbl basis.

Examples. (a) In  $\mathbb{R}^d$ , rational boxes form a basis, so  $\mathbb{R}^d$  is 2<sup>nd</sup> ctbl.  
(b) For a ctbl  $A \neq \emptyset$ , the space  $A^{\mathbb{N}}$  is 2<sup>nd</sup> ctbl because the cylinders form a ctbl basis for  $A^{\mathbb{N}}$ .

Observation. If  $\mathcal{C}$  is a ctbl basis for a metric space  $X$ , the  $\langle \mathcal{C} \rangle_{\sigma} = \mathcal{B}(X)$ .

Prop. For metric spaces, 2<sup>nd</sup> ctblity is equivalent to separability.

Proof. HW

Def. A **measurable space** is a pair  $(X, \mathcal{S})$  where  $X$  is a set and  $\mathcal{S}$  is a  $\sigma$ -algebra on  $X$ .

## Measures.

Def. For a set  $X$  and an algebra  $\mathcal{A}$  on  $X$ , a function  $\mu: \mathcal{A} \rightarrow [0, \infty]$  is said to be:

• **finitely additive** if  $\mu(\bigsqcup_{i \in n} A_i) = \sum_{i \in n} \mu(A_i)$  for all disjoint  $A_0, \dots, A_n \in \mathcal{A}$ .

• **ctbly additive** if  $\mu(\bigsqcup_{i \in \mathbb{N}} A_i) = \sum_{i \in \mathbb{N}} \mu(A_i)$  for all disjoint  $A_0, A_1, \dots \in \mathcal{A}$  with  $\bigsqcup_{i \in \mathbb{N}} A_i \in \mathcal{A}$ .

Def. For a measurable space  $(X, \mathcal{S})$ , a **measure** on  $X$  is a ctblly additive function  $\mu: \mathcal{S} \rightarrow [0, \infty]$  such that  $\mu(\emptyset) = 0$ .

Caution. There is a term finitely additive measure which means a function  $\mu: \mathcal{A} \rightarrow [0, \infty]$  on an algebra  $\mathcal{A}$  that is finitely additive and  $\mu(\emptyset) = 0$ . But finitely additive measures are typically not measures even if  $\mathcal{A}$  is a  $\sigma$ -algebra.


Def. A measure  $\mu$  on a measurable space  $(X, \mathcal{S})$  is called:

- finite if  $\mu(X) < \infty$ .
- probability if  $\mu(X) = 1$ .
- $\sigma$ -finite if  $X$  can be partitioned into ctly many sets from  $\mathcal{S}$  each of which having finite measure.

Observation. (a) A ctly weighted sum of measures is also a measure, i.e. if the  $\mu_n$  are measures on a measurable space  $(X, \mathcal{S})$  and the  $c_n$  are non-negative reals then  $\sum_{n \in \mathbb{N}} c_n \mu_n$  is also a measure on  $(X, \mathcal{S})$ .

(b) A ctly convex combination of probability measures is also a probability measure, i.e. if the  $\mu_n$  are probability measures on a measurable space  $(X, \mathcal{S})$  and the  $c_n$  are non-negative reals with  $\sum_{n \in \mathbb{N}} c_n = 1$ , then  $\sum_{n \in \mathbb{N}} c_n \mu_n$  is also a probability measure on  $(X, \mathcal{S})$ .

Examples. (a) The zero measure  $\mu_0 \equiv 0$  on any measurable space  $(X, \mathcal{S})$ .

(b) The Dirac measure at a point  $x \in X$  is the measure  $\delta_x$  on  $(X, \mathcal{P}(X))$  defined by  $\delta_x(A) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{otherwise} \end{cases}$ . 

(c) The counting measure on  $X$  is the measure  $\mu_c$  on  $(X, \mathcal{P}(X))$  defined by  $\mu_c(A) := \begin{cases} |A| & \text{if } A \text{ is finite} \\ \infty & \text{otherwise} \end{cases}$ .

Note that if  $X$  is ctly, then  $\mu_c = \sum_{x \in X} \delta_x$ .

(d) Given a set  $X$ , define a measure  $\mu$  on the  $\sigma$ -algebra of ctbl/co-ctbl sets:  
$$\mu(A) := \begin{cases} 1 & \text{if } A \text{ is unctbl} \\ 0 & \text{if } A \text{ is ctbl} \end{cases}$$
 This is a measure due to the fact that ctbl unions of ctbl sets are ctbl. Checking this is left as an **exercise**.

Def. Let  $(X, \mathcal{S})$  be a measurable space containing the singleton, i.e.  $\{x\} \in \mathcal{S}$  for all  $x \in X$ .

A measure  $\mu$  on  $(X, \mathcal{S})$  is called

- **atomic** if  $\mu(x) := \mu(\{x\}) > 0$  for some  $x \in X$ . Points of positive measure are called **atoms** of  $\mu$ .
- **purely atomic** if it is a ctbl weighted sum of Dirac measures on  $X$ .
- **nonatomic or atomless** if not atomic, i.e.  $\mu(x) = 0$  for all  $x \in X$ .